

## Today's Stats & Immunizations

Name: \_\_\_\_\_ Date: \_\_\_\_\_

Weight \_\_\_\_\_ / \_\_\_\_\_ % Head Circumference \_\_\_\_\_ / \_\_\_\_\_ %

Length (Height) \_\_\_\_\_ / \_\_\_\_\_ %

Combination (DTaP, IPV, Hib) \_\_\_\_\_ PCV \_\_\_\_\_ Rotavirus \_\_\_\_\_ Flu \_\_\_\_\_ Hepatitis B \_\_\_\_\_ Other: \_\_\_\_\_

Notes: \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

Follow-up Appt: \_\_\_\_\_

Next Well Child Appt: \_\_\_\_\_  
*(9 Months Old)*

**Next Immunizations:**  
*Flu, Anemia Screening, COVID (as eligible)*

## HELPFUL TIPS FOR THIS STAGE OF LIFE:

### Development

- Your laughing, screeching, babbling baby is now starting to take the lead. They are reaching for and moving objects from hand to hand or to their mouth. Your baby is rolling over, trying to sit without support, standing with support, and making sounds to communicate with you.
- Let your baby explore with just enough help to allow her to build confidence. Copying their movements and giving words in response to your baby's sounds lets them know you understand.
- Providing words for feelings, objects that they are interested in, and reading to them helps your baby learn language. Babies also learn through repetition with games (peek-a-boo, patty-cake, etc.) stories, songs, and nursery rhymes.
- Your baby may protest when separated from you and may be starting to seek more comfort from you when around strangers. Consider your own reactions around "strangers" and during times of separation as your baby is taking cues from you.
- Your baby may try self soothing by sucking on a finger or pacifier. Remember that cuddling, holding, and providing nurturing and responsive care does not spoil them.
- Face-to-face interaction with caregivers and avoiding screen time, except for live video chatting with loved ones, will enhance your child's social and verbal development.

### Important Resources

- **Poison Control Center:** (800) 222-1222
- **Suicide and Crisis Lifeline:** 988

### Family Well-being

- Caring for an infant can be exhausting and isolating. It is OK to take time for yourself.
- Consider letting your baby stay with someone you trust in order to reconnect with your partner/friends.

### Infant Care

- Teething rings, a cool washcloth, or a clean finger can be used to massage baby's gums during times of teething discomfort. Avoid teething products with benzocaine.
- Brush teeth with fluoride toothpaste the size of a grain of rice on a soft toothbrush or clean finger.
- Acetaminophen and ibuprofen can help with fever and signs of discomfort.
- It is safe to take your baby outdoors if they are dressed appropriately and protected from the weather. Be cautious about exposing your baby to crowds or to people who may be sick.
- Use sunscreen (SPF 30 or higher).
- Saline drops and suction can be used for nasal congestion, which is common in infants.

### Safety & Child-proofing

- Follow safe sleep guidelines. Place your baby to sleep on their back, on a firm surface, and in their own sleep space. Do not use pillows, blankets, or sleep positioners, as they are not safe for babies.
- Dangling cords, hot liquids, tablecloths, appliances, plastic bags, electrical outlets, and stairs can all be dangerous to an exploring/curious baby.
- Keep household cleaners, vitamins, and medicines out of reach.
- Keep your baby safe with constant supervision during bath time and when on high surfaces such as changing tables, beds, and sofas.
- Avoid walkers. They can tip over and cause your baby harm.
- Be careful of small objects and powder: babies can choke on them.
- Install the car seat in the back seat, rear-facing.
- Make sure smoke and carbon monoxide detectors are working.
- Avoid smoke/tobacco/cigarette/vaping liquid or smoke/marijuana exposure.
- Consider learning infant/child CPR.

## Nutrition

- Breast milk and formula are still the main sources of nutrition and calories.
- Give Vitamin D (400IU/day) to babies receiving breast milk.
- Babies that can sit with support, show good neck/head control, open their mouth for the spoon, and show interest in food are ready to start solids.
- Two or three meal times can be incorporated into your daily routines. You can feed your baby when you and your family are eating. Feed your baby slowly and patiently.
- Follow cues that the baby is hungry or full to help them feel understood and not feel forced to eat.
- Introduce new foods one at a time.
- Mashed and pureed fruits, vegetables, and lentils provide vitamins and minerals. Offer foods naturally rich in iron. (Babies need 11 mg/day of iron.)
- Constipation may increase with the introduction of solids.
- You can also offer a sippy cup and water. Your baby does not need juice, tea, or flavored drinks. **Avoid honey.**
- Only use turmeric/spices from the US to prevent lead poisoning.
- Gassiness is common in early infancy. Try gripe water, simethicone, bicycle legs, or tummy massage for relief. Seek medical advice if inconsolable for 2+ hours.

## Sleep

- Simple, calming, bedtime routines can help prepare your baby for bedtime. Recognize signs that your baby is getting tired in order to know when to slow down and start your routine.
- Your baby's brain is changing and they may be ready for bedtime routines to begin earlier in the evening.
- Many babies are still learning to fall asleep independently. Help your baby learn how to fall asleep on their own by putting them down when they are drowsy, but not fully asleep. Expect a protest if this is a new change for your child.
- Some babies establish a pattern of naps with long stretches of sleep, but others are fine taking shorter naps or napping at less regular times. Recognizing patterns and creating a nap ritual can be helpful. Nap patterns take some trial and error and will change as your baby grows and develops.
- Anticipate sleep disruptions as part of normal development.

## Overseas Travel

If you will be traveling with your family overseas, talk with your doctor about travel questions, or you can request an online pre-travel health consultation through our [eTravel Clinic](#).

| Acetaminophen (Tylenol) Dose Table |            |            |           |            |
|------------------------------------|------------|------------|-----------|------------|
| May repeat every 4 hours.          |            |            |           |            |
| Child's Weight (pounds)            | 6-11 lbs   | 12-17 lbs  | 18-23 lbs | 24-35 lbs  |
| Child's Weight (kilograms)         | 2.7-5.4 kg | 5.5-7.9 kg | 8-10.9 kg | 11-15.9 kg |
| Liquid 160 mg/ 5 milliliters (mL)  | 1.25 mL    | 2.5 mL     | 3.75 mL   | 5 mL       |
| Liquid 160 mg/ 1 teaspoon (tsp)    |            | ½ tsp      | ¾ tsp     | 1 tsp      |

| Ibuprofen (Motrin, Advil) Dose Table                                   |            |           |            |
|--|------------|-----------|------------|
| Do not use in infants under 6 months of age. May repeat every 6 hours. |            |           |            |
| Child's Weight (pounds)  | 12-17 lbs  | 18-23 lbs | 24-35 lbs  |
| Child's Weight (kilograms)   | 5.5-7.9 kg | 8-10.9 kg | 11-15.9 kg |
| Infant Drops 50 mg/ 1.25 mL  | 1.25 mL    | 1.875 mL  | 2.5 mL     |
| Liquid 100 mg/ 5 milliliters (mL)                                      | 2.5 mL     | 3.75 mL   | 5 mL       |
| Liquid 100 mg/ 1 teaspoon (tsp)  | ½ tsp      | ¾ tsp     | 1 tsp      |

## Questions or concerns?



Schedule your next visit on our website.  
[AllegroPediatrics.com](http://AllegroPediatrics.com)



A nurse is available 24 hours a day.  
(425) 827-4600



We have clinics open 7 days a week.